

Nomenclature of the gadolinite supergroup

PETER BAČÍK^{1,*}, RITSURO MIYAWAKI², DANIEL ATENCIO³, FERNANDO CÁMARA^{4,5} and JANA FRIDRICHOVÁ¹

¹ Department of Mineralogy and Petrology, Faculty of Natural Sciences, Comenius University in Bratislava, Mlynská dolina, Ilkovičova 6, 842 15 Bratislava, Slovak Republic

*Corresponding author, e-mail: bacikp@fns.uniba.sk

² Department of Geology and Paleontology, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba 305-0005, Japan

³ Instituto de Geociências, Universidade de São Paulo, Rua do Lago, 562, 05508-080 São Paulo, SP, Brazil

⁴ Dipartimento di Scienze della Terra “Ardito Desio”, Università degli Studi di Milano, via Mangiagalli 34, 20133 Milano, Italy

⁵ CrisDi, Interdepartmental Centre for Crystallography, via Pietro Giuria 5, 10125 Torino, Italy

Abstract: The newly defined gadolinite supergroup approved by the IMA CNMNC (vote 16-A) includes mineral species that have the general chemical formula $A_2MQ_2T_2O_8\varphi_2$ and belong to silicates, phosphates and arsenates. Each site is occupied by: A – Ca, REE (Y and lanthanoids), actinoids, Pb, Mn²⁺, Bi; M – Fe, □ (vacancy), Mg, Mn, Zn, Cu, Al; Q – B, Be, Li; T – Si, P, As, B, Be, S; and φ – O, OH, F. The classification of the gadolinite supergroup is based on the occupancy of A , M , Q , T and φ sites and application of the dominant-valency and dominant-constituent rules. The gadolinite supergroup is divided into two groups defined by prevailing charge occupancy at the T site – Si⁴⁺ in gadolinite group and P⁵⁺ or As⁵⁺ in herderite group. The gadolinite group is divided into the gadolinite and datolite subgroups. The A site is dominantly occupied by divalent cations in the datolite subgroup and by trivalent cations in the gadolinite subgroup. Accordingly, the Q site is dominantly occupied by B³⁺ in the datolite subgroup and by Be²⁺ in the gadolinite subgroup. The herderite group is divided into two subgroups. The herderite subgroup is defined by the dominant divalent cation (usually Ca²⁺) in the A site and Be²⁺ in the Q site, while the M site is vacant. The drugmanite subgroup is defined by the dominance of divalent cations (usually Pb²⁺) in the A site, vacancy in the Q site and the occupation of the M site. Moreover, “bakerite” is discredited as mineral species because it does not meet the conditions of the dominant-constituent rule.

Key-words: gadolinite supergroup; gadolinite; herderite; datolite; drugmanite; nomenclature.

1. Introduction

The newly defined gadolinite supergroup, approved by the Commission on New Minerals, Nomenclature and Classification (CNMNC) of the International Mineralogical Association (IMA) (vote 16-A), includes mineral species that have the general chemical formula $A_2MQ_2T_2O_8\varphi_2$ and belong to silicates, phosphates and arsenates (Table 1). Each site is occupied by (the most common ions or vacancy are underlined):

- A : Ca, REE (Y and lanthanoids), actinoids, Pb, Mn²⁺, Bi;
- M : Fe, □ (vacancy), Mg, Mn, Zn, Cu, Al;
- Q : B, Be, Li;
- T : Si, P, As, B, Be, S;
- φ : O, OH, F.

All minerals of the gadolinite supergroup have monoclinic symmetry and are usually described in the two of three cell choices of the space group no. 14: choice 1, $P2_1/c$ (e.g.

Foit *et al.*, 1973) and choice 3, $P2_1/a$ (e.g. Miyawaki *et al.*, 1984, 1985). For an internally consistent description of all members of the supergroup, we use here only $P2_1/c$.

Historically, the use of the name and classification of the gadolinite supergroup has not been clearly defined; the “group” was referred to as “datolite group” as well as “gadolinite–datolite group”, including or excluding non-silicate members. Following the example of nomenclatures for the apatite (Pasero *et al.*, 2010) and garnet (Grew *et al.*, 2013) supergroups, which included silicate and non-silicate members, the nomenclature of the gadolinite supergroup merges silicate minerals previously belonging to the “gadolinite–datolite group” and phosphates and arsenates with the gadolinite-type structure. Moreover, the need for three-level hierarchies requires the creation of a gadolinite supergroup instead of a gadolinite group.

In the past, there have also been invalid mineral species frequently included in the “datolite group”. Later on, according to further research, they turned out to be transitional between end-member compositions. The list of these invalid species as well as other minerals mistakenly

Table 1. Valid mineral species with gadolinite-type structure.

| | Crystal-chemical formula | | | | | | Lattice parameters | | | | | Ref. |
|--------------------|--------------------------|------------------|-----------------|-----------------|---------------------|-------------------|--------------------|--------------|--------------|-------------|----------------------------|------|
| | <i>A</i> | <i>M</i> | <i>Q</i> | <i>T</i> | <i>O</i> | φ | <i>a</i> (Å) | <i>b</i> (Å) | <i>c</i> (Å) | β (°) | <i>V</i> (Å ³) | |
| Datolite | Ca ₂ | □ | B ₂ | Si ₂ | O ₈ | (OH) ₂ | 4.832 | 7.608 | 9.636 | 90.40 | 354.2 | 1 |
| Homilite | Ca ₂ | Fe ²⁺ | B ₂ | Si ₂ | O ₈ | O ₂ | 4.776 | 7.621 | 9.786 | 90.61 | 356.2 | 2 |
| Gadolinite-(Y) | Y ₂ | Fe ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | 4.768 | 7.565 | 10.000 | 90.31 | 360.7 | 3 |
| Gadolinite-(Ce) | Ce ₂ | Fe ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | 4.820 | 7.580 | 10.010 | 90.28 | 347.1 | 4 |
| Gadolinite-(Nd) | Nd ₂ | Fe ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | 4.822 | 7.699 | 10.136 | 90.23 | 376.2 | 5 |
| Hingganite-(Y) | Y ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | 4.744 | 7.571 | 9.811 | 90.26 | 352.4 | 6 |
| Hingganite-(Ce) | Ce ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | 4.751 | 7.628 | 9.897 | 90.42 | 358.6 | 7 |
| Hingganite-(Yb) | Yb ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | 4.740 | 7.607 | 9.898 | 90.45 | 356.9 | 8 |
| Minasgeraisite-(Y) | Y ₂ | Ca | Be ₂ | Si ₂ | O ₈ | O ₂ | 4.702 | 7.562 | 9.833 | 90.46 | 349.6 | 9 |
| Herderite | Ca ₂ | □ | Be ₂ | P ₂ | O ₈ | F ₂ | 4.763 | 7.677 | 9.745 | 90.67 | 356.3 | 10 |
| Hydroxylherderite | Ca ₂ | □ | Be ₂ | P ₂ | O ₈ | (OH) ₂ | 4.785 | 7.668 | 9.762 | 90.18 | 358.2 | 11 |
| Drugmanite | Pb ₂ | Fe ³⁺ | □ | P ₂ | O ₇ (OH) | (OH) ₂ | 4.643 | 7.986 | 11.111 | 90.41 | 358.2 | 12 |
| Bergslagite | Ca ₂ | □ | Be ₂ | As ₂ | O ₈ | (OH) ₂ | 4.882 | 7.809 | 10.127 | 90.16 | 386.1 | 13 |

References: 1, Foit *et al.* (1973); 2, Miyawaki *et al.* (1985); 3, Miyawaki *et al.* (1984); 4, Segalstad & Larsen (1978); 5, Škoda *et al.* (2016); 6, Demartin *et al.* (2001); 7, Miyawaki *et al.* (2007); 8, Demartin *et al.* (2001); 9, Foord *et al.* (1986); 10, Harlow & Hawthorne (2008); 11, Harlow & Hawthorne (2008); 12, King & Sengier-Roberts (1988); 13, Hansen *et al.* (1984a).

Table 2. Invalid minerals and obsolete names formerly included in the gadolinite supergroup (in *italic*) and minerals shown not to belong to the gadolinite supergroup.

| | Formula | Remark |
|-------------------------------|---|--------|
| <i>Calcybeborosilite</i> -(Y) | (Y,Ca) ₂ (□,Fe ²⁺)(B,Be) ₂ Si ₂ O ₈ (OH,O) ₂ | 1 |
| <i>Calciogadolinite</i> -(Y) | (YCa)Fe ³⁺ Be ₂ Si ₂ O ₈ O ₂ | 2 |
| <i>Yttroceberysite</i> | Y ₂ □Be ₂ Si ₂ O ₈ (OH) ₂ | 3 |
| <i>Xinganite</i> | Y ₂ □Be ₂ Si ₂ O ₈ (OH) ₂ | 4 |
| <i>Nd-dominant hingganite</i> | Nd ₂ □Be ₂ Si ₂ O ₈ (OH) ₂ | 5 |
| <i>Bakerite</i> | Ca ₂ □B ₂ Si _{1.5} B _{0.5} [O _{7.5} (OH) _{0.5}](OH) ₂ | 6 |
| Euclase | BeAlSiO ₄ (OH) | 7 |
| Väyrynenite | MnBePO ₄ (OH) | 8 |

Remarks: 1, name for mineral intermediate in composition between datolite and gadolinite published without approval of CNMNC IMA (Semenov *et al.*, 1963; Povarennykh & Dumatov, 1970); 2, described by Nakai (1938) as ferrous and ferric yttrium calcium beryllium silicate and named according to composition and similarity with gadolinite, but considered questionable, discredited in 2006 as identical with Ca-bearing gadolinite (Burke, 2006). Ito (1967) reported a ferric yttrium calcium beryllium silicate, as the synthetic equivalent of “calciogadolinite”. Ito & Hafner (1974) proposed that (YCa)Fe³⁺Be₂Si₂O₈O₂ is one of the three end-members of a solid solution observed in natural gadolinite samples; 3, obsolete name of hingganite-(Y) (Ding *et al.*, 1981); 4, the pinyin spelling of hingganite-(Y) name (Lulu & Zhi-Zhong, 1986); 5, published composition of hingganite (Pršek *et al.*, 2010) but not yet proposed as mineral species and approved by CNMNC IMA; 6, discredited; 7, first descriptions of the euclase structure indicated similarity to datolite structure but later structure refinement proved differences between both minerals (Biscoe & Warren, 1930); 8, Volborth (1954) suggested the possibility that väyrynenite might be isostructural with herderite. However, structures of väyrynenite and herderite are entirely different as proved by later research (*e.g.* Mrose & von Knorring, 1959).

included in the gadolinite supergroup is reported in Table 2, along with obsolete names of valid end-members. All of them except euclase and väyrynenite (which do not belong to the gadolinite supergroup) are therefore not valid and should be considered discredited from now on.

[100] direction (in the $P2_1/c$ space group). One layer consists of TO_4 and QO_4 tetrahedra, and alternate with layers composed of both $AO_6\varphi_2$ polyhedra and $MO_4\varphi_2$ octahedra (Fig. 1).

2.1. Tetrahedral sites

2. Crystal chemistry of gadolinite-supergroup minerals

The structure of gadolinite-supergroup (GSG) minerals can be described as a layered one composed of two different layers parallel to (100) and alternating along the

Tetrahedral sites in the structure of GSG minerals are linked into regularly alternating 4- and 8-membered rings (Fig. 2a) forming layers (Fig. 2b) (Miyawaki & Nakai, 1996; Cámara *et al.*, 2008). In the $P2_1/c$ setting, all 8-membered rings are stretched along **c**, but rotated 25–30° alternately clockwise and counterclockwise within the

74
75
76
77
7879
8081
82
8384
85
86
87

88

89
90
91
92
93
94




Fig. 1. The structure of gadolinite-super group minerals (structure of gadolinite-(Y) after Cámara *et al.*, 2008): (a) view perpendicular to (100) and (b) view along [010] showing several alternating layers along [100]. Graphics obtained with VESTA 3.1.6 (Momma & Izumi, 2011). (Online version in color.)

(100) plane (Bačík *et al.*, 2014). The TO_4 and QO_4 tetrahedra alternate regularly, every TO_4 tetrahedron is connected to 3 QO_4 tetrahedra and vice-versa.

The TO_4 tetrahedron is occupied by Si, P and As in silicates, phosphates and arsenates, respectively. Charge at the T site can vary from 4^+ to 5^+ , although the observed partial substitution of B at the T site in “bakeritic” datolite (Perchiazzi *et al.*, 2006) implies that its charge can be as low as 3.5^+ . In silicates, significant substitution of P for Si, up to 0.40 atom per formula unit (*apfu*) but usually below 0.05 *apfu*, was observed in GSG minerals from Gemerská Poloma, Bacúch, and Turčok, Slovakia (Uher *et al.*, 2009; Pršek *et al.*, 2010; Bačík *et al.*, 2014), the Jaguaraçu Pegmatite, Minas Gerais, Brazil (Foord *et al.*, 1986), and Skoddefjellet, Svalbard (Majka *et al.*, 2011). In contrast, bergslagite from Tennvatn pegmatite, Norway, contains 0.226 *apfu* Si replacing As and 0.06 *apfu* Si replacing Be (per 4 O + 1 OH formula unit) (Raade *et al.*, 2006). However, based on the current data, it is still unclear whether a continuous solid solution exists between silicates, phosphates and arsenates, or a miscibility gap.

The QO_4 tetrahedra are commonly occupied by B and/or Be. Lithium can be also incorporated in GSG minerals and, if so, likely occurs in the Q and M sites owing to the exchange vectors: (1) ${}^M\text{Fe}^{2+} + {}^A\text{Y}^{3+} \rightarrow {}^M\text{Li}^+ + {}^A(\text{Th}^{4+} + \text{U}^{4+})$ and (2) ${}^Q\text{Be}^{2+} + {}^M\text{Fe}^{2+} \rightarrow {}^Q\text{Li}^+ + {}^M\text{Fe}^{3+}$; the maximum amount of Li allowed in the gadolinite structure is 1.0 *apfu* (Cámara *et al.*, 2008). Drugmanite is unique owing to the vacancy in the Q site (King & Sengier-Roberts, 1988). However, the population of other sites and the structure topology are similar to other GSG minerals,




Fig. 2. Layers of TO_4 (blue) and QO_4 (turquoise) tetrahedra in the structure of GSG minerals (structure of gadolinite-(Y) after Cámara *et al.*, 2008): (a) view perpendicular to (100) and (b) view along [010] showing alternance of several layers along [100]. Graphics obtained with VESTA 3.1.6 (Momma & Izumi, 2011). (Online version in color.)

and thus drugmanite belongs to the GSG. Although H atoms could not be located in its structure, one hydrogen position was tentatively proposed between two O1 sites (see location of O1 anions in Fig. 1), on the basis of the low bond-valence incidence calculated at these anion sites (*ca.* 1.5 valence units (*v.u.*), King & Sengier-Roberts, 1988). It seems therefore that the Q sites are indeed vacant in the structure of drugmanite. Charge at the Q sites can vary from 0 (vacant sites, drugmanite) to 3^+ (datolite).

2.2. The A site

The A site has the highest coordination number (8) in the structure of GSG minerals and forms distorted tetragonal antiprisms (Cámara *et al.*, 2008). The higher coordination number allows the occupancy of the A site by cations with ionic radii between 1.29 (Pb) and 0.977 Å (Lu) (Shannon, 1976). However, Ca (1.12 Å), Y (1.019 Å) and Ce (0.97 Å) are the most common cations occupying the A site. Minor quantities of Mn (0.96 Å) are found at the A site. There is no evidence for the presence of monovalent cations at this site, although in principle there is no limitation in terms of local site dimensions and bond-valence requirements that would exclude the presence of, *e.g.*, Na at the A site. No remarkable deficiency has been reported for the A -site




Fig. 3. Layers of $MO_4\varphi_2$ octahedra (green) and $AO_6\varphi_2$ polyhedra in the structure of GSG minerals (structure of gadolinite-(Y) after Cámara *et al.*, 2008): (a) view along $[100]$ and (b) view along $[010]$. Graphics obtained with VESTA 3.1.6 (Momma & Izumi, 2011). (Online version in color.)

population, which suggests the absence of an *A*-site-vacant end-member. Nevertheless, the charge at the *A* site can vary considerably, from 2^+ to 3^+ .

The $AO_6\varphi_2$ polyhedra are located in layers sandwiched between two tetrahedral layers (Fig. 3a) and also forms a network of irregular 6-membered rings, stretched at angles of *ca.* 20° ($+c$ and $-c$) to the *b* direction. The rings enclose the *M* site, which shares edges with the $AO_6\varphi_2$ polyhedra (Fig. 3b). The $AO_6\varphi_2MO_4\varphi_2$ layer is bounded on both sides by 8-membered rings of tetrahedra (Bačík *et al.*, 2014).

2.3. The *M* site

The distorted $MO_4\varphi_2$ octahedra are bounded on both sides of the $AO_6\varphi_2MO_4\varphi_2$ layer by 4-membered rings of *T* and *Q* tetrahedra. Each *M*-site cation is coordinated to pairs of O2, O4 and φ_5 anions. The O2 and O4 anions connect *TO*₄ and *QO*₄ tetrahedra in neighbouring tetrahedral layers. The φ_5 anions, which are located at the $-a$ and $+a$ apices of *QO*₄ tetrahedra, are shifted out of (001). Thus, $MO_4\varphi_2$ octahedra lie with an O2O4 φ_5 face along (001), with slight deflection of the corners with φ_5 anions. The *M* site is located at the twofold symmetry axis in the corner of the unit cell. Consequently, *M* sites in adjacent layers are located above one another, resulting in tunnels in the *a* direction (Bačík *et al.*, 2014) when the *M* sites are vacant, as in datolite or hingganite-(Y).

The *M* site is commonly occupied by Fe^{2+} as in gadolinites (*e.g.* Miyawaki *et al.*, 1984; Demartin *et al.*, 2001; Cámara *et al.*, 2008) and homilite (Miyawaki *et al.*, 1985), although other divalent cations with ionic radius close to Fe^{2+} (Mg, Mn, Co, Ni, Cu and Zn – Ito & Hafner, 1974; Ni – Foit & Gibbs, 1975) can also enter the site. Nevertheless, the *M* site might also host large divalent cations like Ca in minasgeraisite-(Y), although the local strain requires relaxation and lowering of symmetry (Atencio, unpublished data).

The *M* site might also be occupied by large trivalent cations like Fe^{3+} as suggested by synthetic work (Ito, 1967). This was confirmed by Mössbauer spectroscopy on synthetic “calciogadolinite” (Ito & Hafner, 1974). For the B-free solid-solution series in GSG, a coupled substitution $YFe^{2+}(CaFe^{3+})_{-1}$ is required instead of $YBe(CaB)_{-1}$ (Miyawaki & Nakai, 1996). Ferric iron was suggested for a B-free (SIMS data) and Ca-rich (EMPA data) GSG mineral from Tahara, Japan (Miyawaki *et al.*, 1987) on the basis of $YFe^{2+}(CaFe^{3+})_{-1}$. In contrast, Demartin *et al.* (1993) proposed, on the basis of observed $\langle M-O \rangle$ distances, that iron is entirely in the divalent state in samples from the Alps. However, the incorporation of Li in the *QO*₄ tetrahedron implies that part of the iron might be in the trivalent state (Cámara *et al.*, 2008). On the other hand, the *M* site is commonly vacant as, *e.g.*, in datolite and hingganite (*e.g.*, Foit *et al.*, 1973; Demartin *et al.*, 2001; Miyawaki *et al.*, 2007; Pršek *et al.*, 2010; Majka *et al.*, 2011); the charge is balanced by protonation of two oxygens at the φ_5 anion sites. Both protons are located inside the empty *M* site due to steric hindrance requirements – to avoid repulsion with nearby cations. Therefore, the *M* site is not actually vacant and the $\square(OH)_2Fe_{-1}O_2$ substitution can alternatively be expressed shortly as H_2Fe_{-1} . Charge at the *M* site can vary from 0 (vacant site) to 3^+ .

2.4. Anion sites

There are five anion sites in general positions in the GSG minerals, which account for 10 anions per formula unit. The O2–4 sites are always occupied by oxygen, while O1 may be bonded to one H atom (as in drugmanite, King & Sengier-Roberts, 1988). The φ_5 site hosts O atoms and can host (OH) groups (as in datolite and hingganites) or F (as in herderite). The anion part of gadolinite supergroup (GSG) structure can account therefore for 16–20 negative charges.

3. Definitions and classification principles

The classification of the gadolinite supergroup (Table 3) is based on the occupancy of *A*, *M*, *Q*, *T* and φ sites and on the two rules used for classification of mineral species – dominant-constituent rule and dominant-valency rule (Hatert & Burke, 2008). The dominant-valency rule is used for occupancy of the *A* and *M* site, in which various trivalent and divalent cations (and vacancy in the *M* site) occur, and also for the *T* site, which can be occupied by various tetravalent and pentavalent cations (plus the so far

174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217

218
219
220
221
222
223
224
225
226
227

Table 3. Approved gadolinite-supergroup nomenclature.

| | <i>A</i> | <i>M</i> | <i>Q</i> | <i>T</i> | <i>O</i> | φ | Status |
|---|------------------|------------------|-----------------|-----------------|---------------------|-------------------|--------|
| Gadolinite group (silicates) | | | | | | | |
| <i>Datolite subgroup</i> | A^{2+} | | Q^{3+} | | | | |
| Datolite | Ca ₂ | □ | B ₂ | Si ₂ | O ₈ | (OH) ₂ | A |
| Homilite | Ca ₂ | Fe ²⁺ | B ₂ | Si ₂ | O ₈ | O ₂ | A |
| <i>Gadolinite subgroup</i> | A^{3+} | | Q^{2+} | | | | |
| Gadolinite-(Y) | Y ₂ | Fe ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | A |
| Gadolinite-(Ce) | Ce ₂ | Fe ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | A |
| Gadolinite-(Nd) | Nd ₂ | Fe ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | A |
| Hingganite-(Y) | Y ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | A |
| Hingganite-(Ce) | Ce ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | A |
| Hingganite-(Yb) | Yb ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | A |
| <i>Hingganite-(Nd)^a</i> | Nd ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | H |
| Minasgeraisite-(Y) | Y ₂ | Ca ²⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | A |
| <i>New root name 1^b</i> | Bi ₂ | □ | Be ₂ | Si ₂ | O ₈ | (OH) ₂ | H |
| <i>Calciogadolinite-(Y)^c</i> | CaREE | Fe ³⁺ | Be ₂ | Si ₂ | O ₈ | O ₂ | D |
| <i>New subgroup^d</i> | A^{3+} | | $Q^{2+}Q^+$ | | | | |
| <i>New root name 2^d</i> | REE ₂ | Fe ³⁺ | BeLi | Si ₂ | O ₈ | O ₂ | H |
| <i>New root name 3^d</i> | REE ₂ | Al | BeLi | Si ₂ | O ₈ | O ₂ | H |
| Herderite group (phosphates and arsenates) | | | | | | | |
| <i>Herderite subgroup</i> | A^{2+} | | Q^{2+} | | | | |
| Herderite | Ca ₂ | □ | Be ₂ | P ₂ | O ₈ | F ₂ | A |
| Hydroxylherderite | Ca ₂ | □ | Be ₂ | P ₂ | O ₈ | (OH) ₂ | A |
| Bergslagite | Ca ₂ | □ | Be ₂ | As ₂ | O ₈ | (OH) ₂ | A |
| <i>Fluorbergslagite^e</i> | Ca ₂ | □ | Be ₂ | As ₂ | O ₈ | F ₂ | H |
| <i>Drugmanite subgroup</i> | A^{2+} | | □ | | | | |
| Drugmanite | Pb ₂ | Fe ³⁺ | □ ₂ | P ₂ | O ₇ (OH) | (OH) ₂ | A |
| <i>New root name 4^f</i> | Pb ₂ | Al | □ ₂ | P ₂ | O ₇ (OH) | (OH) ₂ | H |
| <i>New root name 5^g</i> | Pb ₂ | Fe ³⁺ | □ ₂ | As ₂ | O ₇ (OH) | (OH) ₂ | H |
| <i>New root name 6^g</i> | Pb ₂ | Al | □ ₂ | As ₂ | O ₇ (OH) | (OH) ₂ | H |

Status: A, approved; Rd, redefined; Rn, renamed; H, hypothetical end-members based on analytical data or real substitution trends; D, discredited, but hypothetical end-member based on analytical data or real substitution trends.

^a Proposed end-member for Nd-dominant hingganite.

^b Proposed Bi-dominant GSG mineral from Minas Gerais, Brazil described by Foord *et al.* (1986) as Bi-richest composition.

^c Hypothetical end-member for the solid solution observed in natural gadolinite samples suggested by Ito & Hafner (1974) and in natural hingganite samples suggested by Miyawaki *et al.* (1987) and Miyawaki & Nakai (1996). The name of *calciogadolinite-(Y)* was used in published literature but use of a new root name is also possible.

^d Proposed hypothetical subgroup and end-members with shared occupancy of *Q* site by monovalent and divalent cation based on actual LiFe³⁺(BeFe²⁺)₋₁ substitution described in gadolinite-(Y) from Vico, Italy (Cámara *et al.*, 2008) and its Al dominant counterpart.

^e Proposed F-dominant analogue of bergslagite.

^f Proposed end-member based on the actual Al-substitution for Fe³⁺ observed in natural drugmanite (Van Tassel *et al.*, 1979).

^g Arsenate analogue to drugmanite and “*New root name 4*” mineral.

never dominant B³⁺). The gadolinite supergroup is divided into two groups, which are consequently divided into several subgroups.

3.1. Group

Groups within the gadolinite supergroup were named according to the oldest members and are defined by prevailing occupancy at the *T* site on the basis of the dominant-valence rule (Hatert & Burke, 2008). Minerals with dominant Si⁴⁺ belong to the gadolinite group. Minerals with dominant pentavalent ions (P and/or As) belong to the herderite group, also in the case that Si is the most abundant cation at the *T* site but attains less than 1 *apfu* (Fig. 4).

3.2. Subgroup

The classification of a mineral into any of the subgroups is based on the dominant occupancy and valence in *A* and *Q* sites (Table 3). Difficulties in analyzing Be and B suggest that using occupancy of the *Q* sites as a classification criterion is not advisable. Fortunately, determination of the occupation and the dominant charge of the *A* site is usually adequate for assignment of a mineral to a subgroup. However, if the *Q*-site occupancy is not analytically determined and the occupation and the dominant charge of the *A* site are not sufficient for determination of subgroup, the *M*-site occupancy can be used to assign a mineral to the proper subgroup.

228
229
230

231

232
233
234
235
236
237
238
239

240

241
242
243
244
245
246
247
248
249
250
251
252




Fig. 4. Classification diagram of the gadolinite supergroup for determination of groups and subgroups. Inclusion of A^{4+} dominant members would require an additional dimension in the diagram. (Online version in color.)

In the *gadolinite group* there are two subgroups traditionally known – the gadolinite and datolite subgroups. The *A* site is dominantly occupied by divalent cations in the datolite subgroup and by trivalent cations in the gadolinite subgroup. Accordingly, the *Q* site is dominantly occupied by B^{3+} in the datolite subgroup and by Be^{2+} in the gadolinite subgroup.

In the *herderite group* there are two subgroups: 1 – the herderite subgroup is defined by the dominance of divalent cations (usually Ca^{2+}) in the *A* site and Be^{2+} in the *Q* site and 2 – the drugmanite subgroup is defined by the dominance of divalent cations (usually Pb^{2+}) in the *A* site and vacancy in the *Q* site. Although divalent cations occupy the *A* site in both subgroups, no solid solution between subgroups has been reported in natural samples. Therefore, determination of the *A* site occupancy is in general satisfactory for specifying the subgroup (Fig. 4). However, if an intermediate composition is found and if it is impossible to distinguish Be or vacancy at the *Q* site, the occupancy of the *M* site can be used as additional criterion for establishing the proper subgroup in the herderite group.

Any new mineral with different occupancy of the *A* and *Q* sites compared to the above-listed subgroup occupancy (such as the “new root name 2” mineral with mixed occupancy of the *Q* site by Be^{2+} and Li^+ in Table 2, or a hypothetical mineral species with tetravalent cations at the *A* site) will require the definition of a new subgroup.

3.3. Root name, prefixes and suffixes

The root name of GSG minerals is defined by the occupancy of the *A*, *M* and *T* sites.

- Occupancy of the *A* site is expressed in two different ways: (1) if the dominant cation at the *A* site in a new mineral is not a REE, it requires a new root name (*e.g.* datolite with dominant Sr should not be named datolite-

(Sr) or strontio-datolite but should have a different root name) and (2) if a new 4 REE-dominant mineral differs from known species only in the dominant REE, the use of the Levinson-type suffix (Levinson, 1966) of the dominant REE added to the existing root name is recommended (*e.g.* Nd-dominant hingganite should be named *hingganite*-(Nd)). No other use of suffixes is recommended. This also applies to compositions otherwise similar to those of REE-dominant minerals, such as a Bi-dominant analogue of hingganite-(Y) to which a new root name should be given. *Hingganite*-(Bi) is not recommended.

- When the *M* site is occupied (homilite, gadolinite, drugmanite), the dominant cation is usually Fe^{2+} (or Fe^{3+} in drugmanite), no other cation has been found so far to prevail in the *M* site in GSG minerals. If the *M* site occupancy is >0.5 *apfu* and Fe^{2+} (or Fe^{3+} in drugmanite) is not dominant, the use of a new root name is recommended.
- A change in the *T*-site population by heterovalent substitution results in changing the group; consequently, a new root name is necessary. The change in *T*-site population by homovalent substitution (*e.g.* substitution of As for P) does not result in the change of group but the use of a new mineral root name is also recommended, instead of using a prefix. Consequently, the phosphate analogue of drugmanite, if any, should be given a new root name; the use of *phosphdrugmanite* is not recommended.

A prefix could be used to designate an occupant of the $\phi 5$ site by a different anion than in an existing mineral of the supergroup. Currently such a prefix is used only in hydroxylherderite (see below). There is no need to use the prefix oxy- for ϕO dominant datolite-group minerals as its presence causes the change in the *M site occupancy* and subsequently results in a change of root name. Other uses of prefixes are not recommended.

3.4. Adjectival modifiers

Although a subordinate cation does not appear in the name of the GSG minerals, in some cases it may be of sufficient crystal-chemical or petrological interest to deserve mentioning, *e.g.*, 0.70 Sr *apfu* at the *A* site. The International Mineralogical Association Commission on New Minerals and Mineral Names (predecessor of the IMA CNMNC) discredited the use of Schaller’s modifiers, and recommended the use of expressions of the type *Sr-rich* or *Sr-bearing* in such instance (voting proposal 03A; Bayliss *et al.*, 2005). The use of such descriptors is discretionary.

3.5. Definition of mineral species

As stated above, the dominant cations at the *A* and *Q* sites are decisive in the classification of a mineral into a subgroup within the gadolinite supergroup. As the population of the *Q* site is not routinely determined by




Fig. 5. Classification diagram of the gadolinite group for species determination.




Fig. 6. Classification diagram of the herderite subgroup for species determination.

electron microprobe analysis (EMPA), it is appropriate to use only *A* site population for classification of a mineral into a subgroup, whereas populations of the *A*, *M* and *T* sites are critical for defining the species. Similarly, the population of the *M* and *A* (if different from REE) sites is sufficient for determination of the root name. Moreover, in the case of homilite and bergslagite, occupancy of the *M* and *A* sites completely defines the end-member. In contrast, in the gadolinite subgroup it is necessary to determine the dominant trivalent cation in the *A* site for classification into mineral species, thus only the root name is used in the classification diagram (Fig. 5).

Determination of *F* is necessary for classification of herderite and hydroxylherderite (Fig. 6). Drugmanite is the only GSG mineral with Pb in the *A* site and no solid solution with other GSG mineral was observed; only AlFe^{3+}_{-1} substitution (or AlMn^{3+}_{-1} substitution) can produce new end-members of the gadolinite supergroup. Moreover, two As-dominant counterparts can be proposed (Fig. 7).

4. Specific, invalid and discredited gadolinite-supergroup minerals

Hereafter specific, invalid and discredited minerals of the gadolinite supergroup are discussed. We describe minerals with specific crystal-chemical features, minerals of which composition or other analytical data were published but that have not been proposed yet as new mineral species. Moreover, we explain the reasons for discrediting “bakerite”.

4.1. Minasgeraisite-(Y)

Minasgeraisite-(Y) was originally described from the Jaguarçu pegmatite, Minas Gerais, Brazil (Foord *et al.*, 1986), which remains the only known locality. Only

several hundred milligrams are known to exist. Its empirical formula was calculated from the ICP-AES data of bulk sample, namely an admixture of Bi-poor core and Bi-rich rim parts of the heterogeneous material, as $(\text{Y}_{0.72}\text{Ln}_{0.41}\text{Ca}_{0.56}\text{Bi}_{0.31})_{\Sigma 2.00}\text{Ca}_{0.45}\text{Mn}_{0.20}\text{Mg}_{0.08}\text{Fe}_{0.05}\text{Zn}_{0.02}\text{Cu}_{0.01}\square_{0.19})_{\Sigma 1.00}(\text{Be}_{1.55}\text{B}_{0.21}\text{Si}_{0.24})_{\Sigma 2.00}(\text{Si}_{1.95}\text{P}_{0.08})_{\Sigma 2.03}\text{O}_{10}$ on a 10 O basis, where Ln is lanthanoids (La–Lu). As obvious from the formula, the excess in Ca content was assigned to the *M* site. However, no structure-refinement data were obtained from type material to support the presence of Ca at the *M* site. In contrast, there are arguments against it, which must be considered. Calcium has a large ionic radius (1 Å in octahedral coordination – Shannon, 1976) compared to other divalent cations, such as Fe^{2+} . Calcium would lengthen the bonds in the octahedra to at least 2.38 Å, and the $\langle M-\varphi 5 \rangle$ distance would exceed by more than 0.3 Å what has been observed in other structures that have been determined (Bačík *et al.*, 2014). Moreover, the lattice parameters of minasgeraisite-(Y) are similar to those of hingganite-(Y), which suggests that neither the distortion in tetrahedral layers (*b* and *c*), nor the increase in layer distances (*a*) owing the CaFe_{-1} substitution play a role in the minasgeraisite-(Y) structure (Bačík *et al.*, 2014). The $(\text{Na},\text{Mn})\text{O}_6$ octahedron in the structure of nordite-(La) (Bakakin *et al.*, 1970) has an equivalent configuration to the $\text{MO}_4\varphi_2$ octahedra in GSG minerals, between 4-membered Si_4O_{12} rings of SiO_4 tetrahedra. The mean $(\text{Na},\text{Mn})-\text{O}$ interatomic distance, 2.38 Å, suggests that the $\text{MO}_4\varphi_2$ octahedron could accommodate the large Ca^{2+} cation. The interlayer distance of 5.16 Å in nordite-(La) is greater than the interlayer distances in gadolinite and hingganite. Although the structure of the GSG minerals is sufficiently flexible to accommodate a wide range of ionic sizes between the layers, which is one of the reasons gadolinite-subgroup minerals crystallize as both smaller Y- and larger Ce-dominant members, there is no evidence in the cell parameters of minasgeraisite-

379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416




Fig. 7. Classification diagram of the drugmanite subgroup for species determination.

(Y) for an increase of the interlayer distance. Therefore, it is unlikely that the *M* site is dominantly occupied by Ca.

Moreover, calculation of minasgeraisite formula on the basis of 10 anions did not consider the possibility of *M*-site vacancy and OH groups, which results in the overestimation of calculated proportions of the other cations in the formula. Consequently, Ca content was overestimated and could not be accommodated at the *A* site; the excess had to be placed at the *M* site. Recalculation on the basis of $\text{Si} + \text{P} = 2 \text{ apfu}$ (at the *T* site) gave the formula: $(\text{Y}_{0.64}\text{Ln}_{0.36}\text{Ca}_{0.72}\text{Bi}_{0.28})_{\Sigma 2.00}(\text{Ca}_{0.18}\text{Mn}_{0.18}\text{Mg}_{0.07}\text{Fe}_{0.04}\text{Zn}_{0.02}\text{Cu}_{0.01}\square_{0.45})_{\Sigma 1.00}(\text{Be}_{1.80}\text{B}_{0.20})_{\Sigma 2.00}(\text{Si}_{1.92}\text{P}_{0.08})_{\Sigma 2.00}\text{O}_8(\text{OH}_{1.38}\text{O}_{0.62})$. Following the dominant-valence and –constituent rules, the formula indicates that $M = \square$ and $\text{Ca} = \text{Mn}$ in the *M* site. Therefore, the specimen of Foord *et al.* (1986) could be classified as Ca- and Bi-bearing hingganite-(Y) as well as Mn-bearing minasgeraisite-(Y).

Foord *et al.* (1986) reported the chemical data of the Bi-richest portion. The recalculation of the Bi-richest composition on the basis of 2 *T* cations yielded the formula $(\text{Bi}_{0.58}\text{Y}_{0.47}\text{Ln}_{0.29}\text{Ca}_{0.68})_{\Sigma 2.02}(\text{Mn}_{0.23}\text{Mg}_{0.07}\text{Fe}_{0.05}\text{Zn}_{0.02}\text{Cu}_{0.01}\square_{0.62})_{\Sigma 1.00}(\text{Be}_{1.48}\text{B}_{0.52})_{\Sigma 2.00}(\text{Si}_{1.92}\text{P}_{0.08})_{\Sigma 2.00}\text{O}_8(\text{OH}_{1.25}\text{O}_{0.75})$, with the introduction of OH groups according to the procedure of formula calculation recommended for GSG minerals in the next section of this paper. Trivalent cations prevail in the *A* site and Bi is the dominant cation among them, which, according to the dominant-valence rule, is sufficient for the definition of a new mineral species. The simplified formula, $\text{Bi}_2\square\text{Be}_2\text{Si}_2\text{O}_8(\text{OH})_2$, conforms with the general formula of GSG minerals, $A_2MQ_2T_2O_8\varphi_2$.

Recently, the minasgeraisite-(Y) structure was solved and refined in space group *P1* by Cooper & Hawthorne (in preparation), using the chemical composition determined by Foord *et al.* (1986) on material from the same locality. Cooper & Hawthorne (in preparation) found that lowering of the symmetry to acentric triclinic resulted from cation

ordering at the *A* site. In addition, they did not place any Ca at the *M* site and refined the *M*-site occupancy using only the Mn scattering factor. Our nomenclature recommends that, if the occupancy of the *M* site is $>0.5 \text{ apfu}$ and Fe^{2+} is not dominant, a new root name is to be used. However, the “valid” name “minasgeraisite-(Y)” is not applicable for the new root name of $\text{Y}_2\text{MnBe}_2\text{Si}_2\text{O}_{10}$.

Cooper & Hawthorne (in preparation) gave the *M*-site composition $(\square_{1.37}\text{Mn}_{0.63})_{\Sigma 2}$, where both individual *M*-sites in their triclinic model have dominant vacancy. The structural formula $(\text{Y,Ca,Bi,Ln})_2(\square,\text{Mn})(\text{Be,B,Si})_2\text{Si}_2\text{O}_8[(\text{OH},\text{O})_2]$ on the basis of $\text{O} + \varphi = 10$ indicates that the mineral is $(\text{Bi}^{3+},\text{Mn}^{2+})$ -bearing hingganite-(Y). However, Cooper & Hawthorne (in preparation) suggested that ordering at the *A*-site is a valid criterion for recognizing minasgeraisite-(Y) as a mineral species distinct from other GSG minerals, in all of which the *A* site is disordered.

Foord *et al.* (1986) obtained powder XRD data with a Gandolfi camera. However, the chemical composition of the crystal(s) on the camera is not mentioned. The sample was heterogeneous on a very fine scale ($<0.1 \text{ mm}$). Foord *et al.* (1986) reported that rosettes of minasgeraisite are concentrically zoned with medium-purple rims and pale-purple cores. The growth zoning was manifested by light and dark zones. Therefore, an admixture of minasgeraisite-(Y), the Ca- and Bi-bearing variety of hingganite-(Y) and the Bi-analogue of hingganite-(Y) (the new root name 1) was very likely present in the analyzed sample. The discrepancy in unit-cell parameters between Foord *et al.* (1986) and Cooper & Hawthorne (in preparation) suggests a difference in chemical composition between their crystals, although it could also partly result from the different methods used (Gandolfi camera vs. SREF).

Although Cooper & Hawthorne (in preparation) did not confirm the formula for minasgeraisite-(Y) proposed by Foord *et al.* (1986), their formula did retain many of the features of Foord *et al.* (1986) formula. Overall, questions concerning minasgeraisite-(Y) remain open, specifically the number of species present and the implications of ordering at the *A* site. Any redefinition or new-mineral descriptions would require submission of a new proposal to the IMA CNMNC.

4.2. Drugmanite

Drugmanite was first described from Richelle, Belgium (Van Tassel *et al.*, 1979). Electron microprobe data confirmed the presence of Pb, Fe, Al and P, which could produce the possible formulae: $\text{Pb}_{4.02}(\text{Fe}^{3+}_{1.55}\text{Al}_{0.45})\text{P}_{4.00}\text{O}_{17.02}\cdot 3\text{H}_2\text{O}$ or $\text{Pb}_2(\text{Fe}^{3+}_{0.78}\text{Al}_{0.22})(\text{PO}_4)_2(\text{OH})\cdot \text{H}_2\text{O}$. On the basis of powder diffraction pattern, the similarity to gadolinite, datolite–homilite and herderite–hydroxylherderite series, namely GSG, was proposed (Van Tassel *et al.*, 1979). Nothing was known about its structural arrangement until structure refinement, which revealed that drugmanite does indeed have a datolite-like structure (Fig. 8; King & Sengier-Roberts, 1988). However, unlike the other GSG minerals, there is only

455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511




Fig. 8. Comparison of the (a) herderite (Harlow & Hawthorne, 2008) and (b) drugmanite (King & Sengier-Roberts, 1988) structure. Drugmanite contains only TO_4 tetrahedra (dark grey) occupied by P and QO_4 tetrahedra are vacant, whereas herderite has both TO_4 (dark grey) and QO_4 (light grey) tetrahedra occupied by P and Be, respectively. The red (in herderite) and green (in drugmanite) patterns highlight the structural motifs of 8-membered and 4-membered tetrahedral rings similar in both minerals. (Online version in color.)

one type of tetrahedra in the drugmanite structure, which is occupied by P. The QO_4 tetrahedron is vacant. The presence of vacancy in the Q site requires definition of a drugmanite subgroup separately from the herderite subgroup, according to the nomenclature rules.

Similarly to “bakerite”, the presence of molecular H_2O proposed by Van Tassel *et al.* (1979) was not confirmed by King & Sengier-Roberts (1988). However, the reason for the discrepancy between the H_2O content estimated from microprobe data and later structural data in drugmanite and “bakerite” is different. In “bakerite” it was ascribed to impurities but bond-valence requirements indicate that drugmanite contains additional OH group. The additional hydrogen is probably bonded to O1 in TO_4 tetrahedra, thus half of the TO_4 tetrahedra is composed of PO_4 , the other half by PO_3OH (King & Sengier-Roberts, 1988). Moreover, King & Sengier-Roberts (1988) were not able to locate H protons in their structure refinement. Only bond-valence calculations are providing evidence for insufficient charge at $\varphi 5$. However, the presence of cations in the M site suggests that, if H is present at $\varphi 5$, it should be located differently than in other GSG minerals. It could be

allowed by the vacancy in the Q site. Conceivably, drugmanite could be an F-dominant mineral similar to herderite, but Van Tassel *et al.* (1979) did not mention F being detected in their preliminary qualitative analyses by optical spectrography and microchemical tests. Despite all differences, the structure of drugmanite is obviously similar to datolite (Fig. 8). Based on the structural data, the end-member formula of drugmanite was redefined to $Pb_2Fe^{3+}(PO_4)(PO_3OH)(OH)_2$ (King & Sengier-Roberts, 1988). For the present, this formula can be written as $Pb_2Fe^{3+}\square_2P_2[O_7(OH)](OH)_2$, which is consistent with the general formula of the GSG. However, it is still necessary to determine the exact $\varphi 5$ occupancy and H location in the structure.

Chemical analyses of drugmanite original material displayed a significant proportion of Al substituting for Fe^{3+} in the octahedral M site (Van Tassel *et al.*, 1979). The existence of an Al-dominant end-member in the drugmanite subgroup is therefore very likely. Moreover, two arsenates analogous to the herderite subgroup can be proposed (Table 3). Consequently, we prefer the definition of a subgroup with one valid end-member and one hypothetical end-member, instead of considering drugmanite as an unassigned member of the herderite group.

4.3. Hingganite-(Nd)

There is one valid Nd-dominant mineral species in the gadolinite group – gadolinite-(Nd), which was recently described as a new mineral from Malmkärä mine, WSW of Norberg, Sweden (with type specimen in the Moravian Museum, Czech Republic, catalogue number B 11298) and approved by the IMA-CNMNC (IMA No. 2016-013: Škoda *et al.*, 2016). Similarly to gadolinite-(Nd), Nd-rich hingganite from Bacúch (Western Carpathians, Slovakia) displays Nd predominating over Y (Nd/Y = 1.27–1.41) and other lanthanoids (Nd/Ce = 1.88–2.20; Nd/Sm = 2.38–2.67; Nd/Gd = 3.49–4.76, and higher) (Pršek *et al.*, 2010). The Nd content reaches 0.57 *apfu* Nd, and concentrations of Y range from 0.40 to 0.49 *apfu*. In addition, Nd-dominant hingganite is generally Ca-poor (≤ 0.01 *apfu*). Therefore, Nd-rich hingganite from Bacúch (Western Carpathians, Slovakia) corresponds to the new Nd-dominant member of the GSG – “hingganite-(Nd)”. Nonetheless, it should be fully described and submitted to the CNMNC IMA for approval as a valid mineral species.

4.4. “Bakerite” discredited

“Bakerite” was first described by Giles (1903) with the chemical formula $Ca_8B_{10}Si_6O_{30}(OH)_{10} \cdot H_2O$. According to Pemberton (1971), the true locality of Giles’ material is Corkscrew Canyon, Death Valley, Inyo County, California. Palache *et al.* (1951) documented structural relationship between “bakerite” and datolite on powder X-ray diffraction patterns and proposed the chemical formula $Ca_8B_{10}Si_6O_{30}(OH)_{10}$, recalculated on the basis of 40 O and φ . Moreover, in the original formula of “bakerite”, there was molecular H_2O present (Giles, 1903), which

should have been located near the *M* site but was later ascribed to the presence of impurities and omitted from the formula $\text{Ca}_4\Box\text{B}_5\text{Si}_3\text{O}_{15}(\text{OH})_5$ (Perchiazzi *et al.*, 2004). This formula can be written as $\text{Ca}_2\Box\text{B}_2(\text{Si}_{1.5}\text{B}_{0.5})[\text{O}_{7.5}(\text{OH})_{0.5}](\text{OH})_2$, which is consistent with the general formula of GSG minerals.

“Bakerite” is therefore characterized by substitution of B for Si in the *T* site, with a proposed charge-balance by OH placed in the O1 site. Hydrogen bonded to O1 has a different location compared to the H (at *H5* site) bonded to $^{\circ}\text{O}$. It lies within the tetrahedral layer, whereas H5 is located on the edge of the tetrahedral layer. Hydrogen bonding between H and O at O1 of neighbouring TO_4 octahedra could result in the structural arrangement with very well-ordered alternation of B and Si with the ratio of 5:3. As a result, there is supposedly a significant gap between the compositions of “bakerite” and datolite; there are no intermediate compositions between them (Perchiazzi *et al.*, 2004).

However, it is not clear if this applies only to samples studied by Perchiazzi *et al.* (2004) or if it is a universal limit for BSi_{-1} substitution. In this case, the application of the *dominant constituent rule* (Hatert & Burke, 2008) results in the discreditation of “bakerite”. Boron occupies only 25% of the *T* site, is therefore not dominant in this site and “bakerite” should not be treated as end-member. The only possibility to preserve “bakerite” as an end-member with respect to the *dominant-constituent rule* is the case of a highly ordered incorporation of B into one specific *T* site, resulting in the splitting of *T* into two sites with consequent reduction of structural symmetry. In this case B could occupy exactly 50% of the site, which would be consistent with the definition of the end-member. However, this was not fully proven by the structural study (Perchiazzi *et al.*, 2004) and “bakerite”, therefore, is discredited (IMA CNMNC 16-A).

5. Calculation of the formula

5.1. Basis of formula calculation

There are several possible procedures for calculation of chemical formula of GSG minerals. The calculation on the basis of 10 anions is proper only when all chemical data are available. In other cases, any calculation on the basis of anions is wrong in principle owing to the presence of OH groups, the possibility of variable amount of Fe^{3+} , and the contents of Be, B and Li, which cannot be quantified by EMPA (or if they are – as in case of B – the precision of measurement is still unsatisfactory, see Ottolini *et al.*, 2002). Therefore, if only EMPA data are available, calculation on the cation basis is a proper option. Calculation on the basis of 5 cations is inappropriate in principle owing to the presence of the *M* site vacancy. Thus there remains three options of calculation – on the basis of (i) 4 (*A* + *T*), (ii) 2 *A*, and (iii) 2 *T* cations.

Calculation on the basis of 4 (*A* + *T*) cations is not advisable owing to possible CaFe_{-1} and BSi_{-1} or AlSi_{-1} substitutions. We presume that CaFe_{-1} substitution is unlikely due to the structural properties of the $\text{MO}_4\varphi_2$

octahedron, but the calculation on the basis of 2 *T* cations is a convenient test for this presumption. Moreover, it could take into account the possibility of *A*-site vacancy although, until now, there is no analytical evidence for it to be significant. In contrast the substitution of B or Be for Si at the *T* sites could also generate certain non-stoichiometry if the formula is calculated on basis of 4 (*A* + *T*) or 2 *T* cations. In this case, the calculation on the basis of 2 *A* cations yields the proper formula. Substitution of Al for Si is negligible in GSG minerals; Al likely fractionates at the *M* site as documented by analyses of Al-bearing material. If no BSi_{-1} substitution is present, like in most GSG minerals, the recommended practice is to calculate formulae on the basis of 2 *T* and 2 *A* cations simultaneously and subsequently to choose the stoichiometrically more adequate formula, with a preference for the 2 *T*-cation calculation.

5.2. The content of O, OH and F in the \Box site

The population in the φ site depends directly on the occupancy of the *M* site, as hydrogen cations are located in the vacant space of the $\text{MO}_4\varphi_2$ octahedron. Fluorine content can be measured by EMPA. Provided that accuracy on F by means of EMPA is good enough, the OH content can be calculated as twice the *M*-site vacancy minus F and then the O content is equal to $2 - (\text{OH} + \text{F})$. The presence of molecular H_2O , which was presumed to be located in the channels formed by vacant *M* sites in “bakerite”, was not confirmed by later research (Perchiazzi *et al.*, 2004).

5.3. Boron, beryllium and lithium

Beryllium and lithium in the *Q* site cannot be determined by EMPA; they should be calculated or determined by other methods. Special EMPA conditions are required for B determination, such as using a layered dispersion element as monochromator for $\text{BK}\alpha$ line with long *d*-spacing and accurate control of peak area and peak shape factors, as well as crystallographic orientation control (Bastin & Heijligers, 1986a and b, 1991). Moreover, EMPA measurements of B require long counting times (*e.g.*, 60 s on peak, 30 s on background) to improve statistics (McGee & Anovitz, 1996). “Routine” operating conditions (10 kV, 15 nA beam current, 30 s counting time) yielded an approximate detection limit of $\sim 0.2\text{--}0.3$ wt% B_2O_3 (McGee *et al.*, 1991). However, the relative analytical error (1σ) of B measurement on EMPA is still relatively high, at the level of several percent (Kutzschbach *et al.*, 2016). Nevertheless, structure refinement of many GSG minerals revealed occupancy of the *Q* site by B along with Be (Demartin *et al.*, 1993, 2001; Cámara *et al.*, 2008). Thus, the sum of Be and B should be specifically mentioned in tabulations of EMPA data even if B and Be were not analyzed or calculated.

Moreover, lithium can be an important constituent of GSG minerals, and its presence should be systematically checked. Lithium should be suspected when (i) the

645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698

(Th + U) content is significant or (ii) the sum Si + B + Be is lower than 2 *apfu*. Hence, metamict samples (containing Th and U) could have significant Li contents (Cámara *et al.*, 2008).

Provided that there is no Li in GSG minerals, the calculation of approximate B content may be done, if we assume that all B enters the structure through the CaB (REE)₋₁Be₋₁ substitution, if Fe(Mn,Mg,Al) = 0 *apfu*. Subsequently, B would be proportional to the Ca content.

Another approach in B/Be ratio calculation is based on the presumption that all Fe or Mn in the mineral is in divalent form and all ⁹OH serves to balance vacancy in the M site. In this case, Be and B can be calculated from charge balance according to $Be = 6 - [20 - \sum PC - OH]$ (where $\sum PC$ is the sum of charges for all cations except B and Be) and $B = 2 - Be$. However, this procedure can be used only if data for all measurable cations are available and a correct calculation producing stoichiometrically more adequate formula is chosen.

Alternatively, if structural data are available, the Q site population in terms of Be and B can be obtained using the calibration by Demartin *et al.* (2001) based on $\langle Q-O \rangle$ distances, corrected by Cámara *et al.* (2008) for the possible presence of Li.

Acknowledgements: We are indebted to Frédéric Hatert and all members of the IMA Commission on New Minerals, Nomenclature and Classification for their helpful suggestions and comments. We also thank Sergey V. Krivovichev for editorial handling and Edward Grew and Milan Novák for their detailed reviews.

Appendices

Appendix Gadolinite-supergroup end-members: species, end-member formulae, etymology, type localities and eventual location of holotypes

All known GSG minerals are monoclinic $P2_1/c$. Representative chemical analyses are listed in Table S1, freely available online as Supplementary Material linked to this article on the GSW website of the journal: <http://eurjmin.geoscienceworld.org>.

Datolite

Structural formula: Ca₂□B₂Si₂O₈(OH)₂.

IMA number: Grandfathered (1806).

Description: Datolite is the second described GSG mineral; Klaproth (1806) found it in Nødebros Mine, Norway. Its formation is commonly connected with volcanic or hydrothermal processes (Giles, 1903; Pemberton, 1971; Bellatreccia *et al.*, 2006; Zaccarini *et al.*, 2008). Datolite previously described as “bakerite” is also a microcrystalline material in various genetic environments, including diabase spilites (Baysal & Dilekoz, 1975),

diopside-bearing marble (Sabina, 1978), and gehlenite–spurrite-bearing skarns (Kusachi *et al.*, 1994). It is usually colorless or white, commonly with a greenish tinge; maybe grayish, yellow, green, red, pink. Datolite usually contains only insignificant proportion of other elements; the Fe substitution is limited up to 0.16 Fe *apfu*, other elements are in even lower proportions (Grew, 1996, and references therein). Excess in B and its substitution for Si in the T site is characteristic for “bakerite” (Perchiazzi *et al.*, 2004).

Etymology: Named from the Greek, meaning *to divide*, in reference to the granular texture of the massive variety.

Type locality: Nødebros Mine (Nøddebro), Arendal Iron Mines, Arendal, Aust-Agder, Norway.

Selected references: Klaproth (1806), Foit *et al.* (1973), Bellatreccia *et al.* (2006), and Rinaldi *et al.* (2010).

Homilite

Structural formula: Ca₂Fe²⁺B₂Si₂O₈O₂.

IMA number: Grandfathered (1876).

Description: Homilite was first described by Paijkull (1876). Structure was solved by Miyawaki *et al.* (1985) on the material from type locality. The composition of the described homilite is near end-member with only up to 0.07 *apfu* ⁹□ and 0.03 *apfu* Mn (Miyawaki *et al.*, 1985). Homilite usually occurs in pegmatite bodies (Paijkull, 1876; Brögger, 1890). It was also discovered in a volcanic ejectum at Tre Croci, Vico volcanic complex, Viterbo Province, Latium, Italy (Pierini, 2004; Boiocchi *et al.*, 2006). A possible third locality was reported in Romania (Hirtopanu *et al.*, 2003).

Etymology: Name from the Greek meaning *to occur together*, in allusion to its association with meliphanite and allanite.

Type locality: Stokø Island, Langesundfjord, Norway.

Location of holotype: Muséum Nationale d’Histoire Naturelle, Galerie Nationale de Minéralogie et de Géologie, Paris, France (catalogue # 78142), Naturhistoriske Riksmuseet, Sektionen för Mineralogi, Stockholm, Sweden (catalogue # 531010).

Selected references: Paijkull (1876) and Miyawaki *et al.* (1985).

Gadolinite-(Y)

Structural formula: Y₂Fe²⁺Be₂Si₂O₈O₂.

IMA status: Renamed (1987).

Description: Gadolinite-(Y) has been an important mineral in the history of the discovery of rare-earth elements since yttria was first separated by J. Gadolin in 1794. The structure of gadolinite-(Y) was derived from datolite structure by simple replacement of atoms (Ito & Mori, 1953). Afterwards, the structure was refined by Miyawaki *et al.* (1984). Gadolinite-(Y) forms accessory phases in REE- and Be-rich granites and pegmatites, and also in some metamorphic rocks (Demartin *et al.*, 1993; Holtstam & Andersson, 2007). Recent classification of rare-element granitic pegmatites (Černý, 1991; Černý & Ercit, 2005) includes the gadolinite NYF (Nb-Y-F) sub-type family with typical mineralogical association consisting of

746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787

gadolinite, fergusonite, euxenite, topaz and beryl. Most enriched in Ca (up to 0.84 *apfu*) is gadolinite-(Y) from Baveno, Italy (Pezzotta *et al.*, 1999), Skodefjellet, Svalbard (Majka *et al.*, 2011), and Dlhá Dolina, Slovakia (Bačík *et al.*, 2014). The highest contents of Mg in gadolinite-(Y) (up to 0.048 *apfu*) were found in Kola Peninsula, Russia (Voloshin *et al.*, 2002), Michalowice, Poland (Kozłowski & Dzierżanowski, 2007) and Mn (up to 0.048 *apfu*) in Skodefjellet, Svalbard (Majka *et al.*, 2011), Michalowice, Poland (Kozłowski & Dzierżanowski, 2007). Aluminium-rich gadolinite (up to 0.19 *apfu*) occurs in Kola Peninsula, Russia (Voloshin *et al.*, 2002), Rode Ranch, Texas, USA (Gibson & Ehlmann, 1970). Gadolinites-(Y) specially enriched in lanthanoids are from Dlhá Dolina, Slovakia – Dy 0.152 *apfu*, Er 0.079 *apfu* (Bačík *et al.*, 2014), Baveno, Italy – Nd 0.271 *apfu*, Gd 0.147 *apfu*, Dy 0.108 *apfu* (Pezzotta *et al.*, 1999), and Turčok, and Slovakia – Nd 0.165 *apfu*, Dy 0.144 *apfu*, Er 0.089 *apfu* (Uher *et al.*, 2009; Bačík *et al.*, 2014).

Etymology: Named after the Finnish chemist Johan Gadolin (1760–1852), who discovered yttrium.

Type locality: Ytterby, Sweden.

Modifications of end-member definition: Renamed after CNMNC IMA approval owing to description of gadolinite-(Ce).

Selected references: Klaproth (1802) and Miyawaki *et al.* (1984).

Gadolinite-(Ce)

Structural formula: $\text{Ce}_2\text{Fe}^{2+}\text{Be}_2\text{Si}_2\text{O}_8\text{O}_2$.

IMA number: 1987 s.p.

Description: Gadolinite-(Ce) is similar to gadolinite-(Y), but is lanthanoid-dominant with most abundant Ce (Segalstad & Larsen, 1978). It is commonly rich in Mg (up to 0.191 *apfu* and Mn (up to 0.148 *apfu*) in several localities in Sweden (Holtstam & Andersson, 2007). Variable Y content (up to 0.821 *apfu*) is observable in Baveno, Italy, gadolinite (Pezzotta *et al.*, 1999). Calcium-rich gadolinites were found in Cuasso al Monte, Italy (up to 0.263 *apfu*) (Pezzotta *et al.*, 1999) and Skien in Norway (up to 0.283 *apfu*) (Segalstad & Larsen, 1978).

Etymology: Named for its cerium content and the relationship to gadolinite-(Y).

Type locality: Bjørkendalen region, and in the Bakken quarry, Tvedalen, Norway.

Location of holotype: Mineralogical-Geological Museum, University of Oslo, Oslo, Norway (catalogue # 21325).

Selected reference: Segalstad & Larsen (1978).

Gadolinite-(Nd)

Structural formula: $\text{Nd}_2\text{Fe}^{2+}\text{Be}_2\text{Si}_2\text{O}_8\text{O}_2$.

IMA number: 2016-013.

Description: Gadolinite-(Nd) is similar to gadolinite-(Y), but is lanthanoid-dominant with most abundant Nd (Škoda *et al.*, 1978).

Etymology: Named for its neodymium content and the relationship to gadolinite-(Y).

Type locality: Malmkärra mine, ca. 2.5 km WSW of Norberg, Sweden.

Location of holotype: Moravian Museum, Brno, Czech Republic (catalogue # B 11298).

Selected reference: Škoda *et al.* (2016).

Hingganite-(Y)

Structural formula: $\text{Y}_2\text{Be}_2\text{Si}_2\text{O}_8(\text{OH})_2$.

IMA number: Renamed (1981-052).

Description: Hingganite-(Y) is the hydrous, Fe-free analogue of gadolinite-(Y). The most Fe-rich hingganite-(Y) is from Cuasso al Monte, Italy (up to 0.496 *apfu*) (Demartin *et al.*, 1993). The most Ca-rich hingganite-(Y) (up to 0.961 *apfu*) is from Dara-i-Pioz Glacier, Tajikistan (Pekov *et al.*, 2000), then from Vico lake, Italy (Cámara *et al.*, 2008), Vlastějovice, Czech Republic (Bačík *et al.*, 2014), Skodefjellet, Svalbard (Majka *et al.*, 2011), Turčok, Slovakia (Uher *et al.*, 2009), Gemerská Poloma, Slovakia (Bačík *et al.*, 2014), Mont Blanc, France (Demartin *et al.*, 1993), and Strange Lake, Canada (Jambor *et al.*, 1998). Specimens rich in Mn (up to 0.101 *apfu*) are from Vlastějovice, Czech Republic (Bačík *et al.*, 2014) and Skodefjellet, Svalbard (Majka *et al.*, 2011). The specimens from Ploskaya Mt, Keivy Mountains, Kola Peninsula, Russia (Voloshin *et al.*, 2002) are rich in lanthanoids – Yb (up to 0.393 *apfu*), Dy (up to 0.223 *apfu*), and Er (up to 0.229 *apfu*).

Etymology: Named after its type locality.

Type locality: Heilongjiang, and greater Xingan (Hinggan) area, Manchuria, China.

Location of holotype: National Museum for Geology, Beijing, China.

Modifications of end-member definition: Renamed after description of hingganite-(Yb).

Selected references: Ding *et al.* (1984) and Demartin *et al.* (2001).

Hingganite-(Ce)

Structural formula: $\text{Ce}_2\text{Be}_2\text{Si}_2\text{O}_8(\text{OH})_2$.

IMA number: 2004-004.

Description: Hingganite-(Ce) is the Ce analogue of hingganite-(Y) (Ding *et al.*, 1984), and can be also expressed as the hydrated Fe-free analogue of gadolinite-(Ce) (Segalstad & Larsen, 1978). Hingganite-(Ce) richest in Fe (up to 0.486 *apfu*) is from Bacúch, Slovakia (Pršek *et al.*, 2010) and Strange Lake, Canada (Jambor *et al.*, 1998); in Y (up to 0.936 *apfu*) from Bacúch, Slovakia (Pršek *et al.*, 2010); in Mn (up to 0.024 *apfu*) from Strange Lake, Canada (Jambor *et al.*, 1998). Hingganites-(Ce) significantly enriched in lanthanoids are from Bacúch, Slovakia with Ce 0.353 *apfu*, Gd 0.149 *apfu* and in some compositions with Nd up to 0.566 *apfu* which corresponds to hingganite-(Nd) composition (Pršek *et al.*, 2010), and from Tahara, Japan – Ce 0.814 *apfu*, La 0.322 *apfu* (Miyawaki *et al.*, 2007).

Etymology: Named for its relationship to hingganite-(Y).

Type locality: Tahara, Nakatsugawa, Gifu Prefecture, Japan.

Location of holotype: National Museum of Nature and Science, Japan (catalogue # NSM-M28552).

Selected reference: Miyawaki *et al.* (2007).

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

Hingganite-(Yb)

Structural formula: $\text{Yb}_2\text{Be}_2\text{Si}_2\text{O}_8(\text{OH})_2$.

IMA number: 1982-041.

Description: Hingganite-(Yb) is the Yb analogue of hingganite-(Y). The best known hingganite-(Yb) is from Ploskaya Mt, Keivy Mountains, Kola Peninsula, Russia. It is characterized by extremely low amount of Fe (up to 0.006 *apfu*) and conversely high amounts of lanthanoids and Y, especially Yb (up to 0.940 *apfu*) and Y (up to 0.832 *apfu*) (Voloshin *et al.*, 2002).

Etymology: Named for its relationship to hingganite-(Y).

Type locality: Mt. Ploskaya, Keivy massif, Kola Peninsula, Russia.

Location of holotype: Fersman Mineralogical Museum, Moscow, Russia (catalogue # 84278-80), Institute of Geology, Kola Science Center of the Russian Academy of Sciences, Apatity, Russia (catalogue # 5768), and Mining Museum, St.Petersburg, Russia (catalogue # 1590/1).

Selected references: Voloshin *et al.* (1983) and Yakubovich *et al.* (1983).

Minasgeraisite-(Y)

Structural formula: $\text{Y}_2\text{CaBe}_2\text{Si}_2\text{O}_8\text{O}_2$.

IMA number: 1983-090.

Description: Minasgeraisite is solely known from the type locality and contains major Y (0.72 *apfu*) and lanthanoids (sum of 0.41 *apfu*), Bi (0.31 *apfu*), Ca (0.56 *apfu* in *A* site and 0.45 *apfu* in *M* site, see discussion in text) and Mn (up to 0.20 *apfu*) and very low Fe (0.05 *apfu*). It occurs as a sparse, accessory, late-stage mineral in small druses in a zoned, complex granitic pegmatite. Minasgeraisite coats and is intergrown with milarite, muscovite, quartz, albite (Foord *et al.*, 1986).

Etymology: Named after its locality.

Type locality: The Jaguaracu Pegmatite, Minas Gerais, Brazil, also known as the Jose Miranda mine or the Zé Pinto or José Pinto mine.

Location of holotype: U.S. National Museum, Washington, USA (catalogue # 164209).

Selected reference: Foord *et al.* (1986).

Herderite

Structural formula: $\text{Ca}_2\text{Be}_2\text{P}_2\text{O}_8\text{F}_2$.

IMA number: Grandfathered (1828).

Description: Herderite is ideally the F end-member in a solid-solution series with hydroxyl-herderite and occurs in complex granite pegmatite (Harlow & Hawthorne, 2008). Herderite from Mogok Myanmar is rich in F (0.748 *apfu*) and Na (0.015 *apfu*). Sodium is the main additional constituent in the Mogok sample (Harlow & Hawthorne, 2008).

Etymology: Named for Siegmund August Wolfgang von Herder (1776–1838), mining official in Freiberg, Saxony, Germany.

Type locality: Ehrenfriedersdorf, Saxony, Germany.

Location of holotype: Werner-Sammlung der Bergakademie Freiberg (Massanek *et al.*, 1999) and may be considered a topotype (Embrey & Hey, 1970).

End-member definition: The F-dominant member in the herderite subgroup.

Selected references: Haidinger (1828) and Harlow & Hawthorne (2008).

Hydroxylherderite

Structural formula: $\text{Ca}_2\text{Be}_2\text{P}_2\text{O}_8(\text{OH})_2$.

IMA number: Renamed (2007) s.p.

Description: Hydroxyl-herderite occurs in granite pegmatites, greisens, and miarolitic cavities associated with granites and granitic pegmatites (Černý, 2002; Grew, 2002). Hydroxylherderite from Ehrenfriedersdorf Germany has almost equal contents of F (0.483 *apfu*) and OH (0.517 *apfu*) (Harlow & Hawthorne, 2008) but hydroxylherderite from Manitoba Canada is much closer to end-member, with up to 0.996 OH *apfu*, low amount of Cl (up to 0.004 *apfu*), F (up to 0.007 *apfu*) and some Na (up to 0.021 *apfu*) (Černá *et al.*, 2002).

Etymology: Named for the composition and relationship to herderite.

Type locality: Paris, Maine, USA.

End-member definition: The (OH)-dominant member in the herderite subgroup.

Selected references: Penfield (1894), Černá *et al.* (2002), and Harlow & Hawthorne (2008).

Drugmanite

Structural formula: $\text{Pb}_2\text{Fe}^{3+}\text{P}_2[\text{O}_7(\text{OH})](\text{OH})_2$.

IMA number: 1978-081.

Description: Drugmanite is a very rare mineral, formed at low temperatures in vugs in mineralized limestones, an oxidation product of disseminated sulfides. It contains up to 0.223 Al *apfu* (Van Tassel *et al.*, 1979). It was found at the type locality Richelle, Belgium (Van Tassel *et al.*, 1979) and Neue Hoffnung Mine, Eifel, Germany (Schnorrer & Schäfer, 1999).

Etymology: Named for Julien Drugman (1875–1950), Belgian mineralogist.

Type locality: Richelle, near Visé, Belgium.

Location of holotype: Belgian Royal Institute of Natural Sciences, Brussels, RN5210; University of Liège, Liège, Belgium, 19347.

Selected references: Van Tassel *et al.* (1979) and King & Sengier-Roberts (1988).

Bergslagite

Structural formula: $\text{Ca}_2\text{Be}_2\text{As}_2\text{O}_8(\text{OH})_2$.

IMA number: 1983-021.

Description: Bergslagite occurs in thin veins in pieces of hematite ore found on mine dumps from a metamorphosed Fe–Mn orebody in Långban, Sweden. It is associated with manganian diopside, manganberzeliite, tilasite, svabite, hematite, calcite, barite (Långban, Sweden); hematite, muscovite (Sailauf, Germany). No other elements are present in significant proportion except Si substituting for As up to 0.08 *apfu* (Hansen *et al.*, 1984a and b). It was described at several other localities. In Tennvatn pegmatite, Norway, bergslagite contains 0.226 *apfu* Si replacing As and 0.06 *apfu* Si replacing Be (per 4 O + 1 OH formula unit, Raade *et al.*, 2006).

Etymology: Named for the occurrence at Långban, in the Bergslagen region of Sweden.

Type locality: Långban, Värmland, Sweden.

Location of holotype: University of Copenhagen, Copenhagen, Denmark; National Museum of Natural History, Washington, DC, USA, 162582.

Selected references: Hansen *et al.* (1984a, b).

Q2 Uncited references

Leavens *et al.* (1978) and Uher *et al.* (2010).

References

- Bačík, P., Fridrichová, J., Uher, P., Pršek, J., Ondrejka, M. (2014): Crystal chemistry of gadolinite-datolite group minerals. *Can. Mineral.*, **52**, 625–642.
- Bakakin, V.V., Belov, N.V., Borisov, S.V., Solovieva, L.P. (1970): The crystal structure of nordite and its relationships to melilite and datolite-gadolinite. *Am. Mineral.*, **55**, 1167–1181.
- Bastin, G.F. & Heijligers, H.J.M. (1986a): Quantitative electron probe microanalysis of boron in some binary borides. in “Microbeam analysis”, A.D. Romig Jr. & W.F. Chambers, eds. San Francisco Press, Inc., California, 285–288.
- Bastin, G.F. & Heijligers, H.J.M. (1986b): Quantitative electron probe microanalysis of boron in binary borides. Internal Report Eindhoven University of Technology, Eindhoven, the Netherlands.
- Bastin, G.F. & Heijligers, H.J.M. (1991): Quantitative electron probe microanalysis of ultra-light elements (boron-oxygen). in “Electron probe quantification”, K.F.J. Heinrich & D.E. Newbury, eds. Plenum Press, New York, USA, 145–161.
- Bayliss, P., Kaesz, H.D., Nickel, E.H. (2005): The use of chemical-element adjectival modifiers in mineral nomenclature. *Can. Mineral.*, **43**, 1429–1433.
- Baysal, O. & Dilekoz, E. (1975): A study of bakerite. *Bull. Miner. Res. Explor. Inst. Turkey*, **84**, 90–96.
- Bellatreccia, F., Della Ventura, G., Cámara, F. (2006): Datolite: a new occurrence in volcanic ejecta (Pitigliano, Toscana, Italy) and crystal-structure refinement. *Rend. Fis. Acc. Lincei*, **17**, 289–298.
- Biscoe, J. & Warren, B.E. (1930): The structure of euclase HBeAlSiO_5 . *Z. Kristallogr.*, **86**, 292–297.
- Boiocchi, M., Callegari, A., Ottolini, L. (2006): The crystal structure of piergorite-(Ce), $\text{Ca}_8\text{Ce}_2(\text{Al}_{0.5}\text{Fe}^{3+}_{0.5})_{\Sigma 1}(\square, \text{Li}, \text{Be})_2\text{Si}_6\text{B}_8\text{O}_{36}(\text{OH}, \text{F})_2$: a new borosilicate from Vetralla, Italy, with a modified hellandite-type chain. *Am. Mineral.*, **91**, 1170–1177.
- Brögger, W.C. (1890): Die Mineralien der Syenitpegmatitgänge der südnorwegischen Augit- und Nephelinsyenite. *Z. Kristallogr. Mineral. B (Leipzig)*, **16**, 1–663.
- Burke, E.A.J. (2006): Mass discreditation of GQN minerals. *Can. Mineral.*, **44**, 1557–1560.
- Cámara, F., Oberti, R., Ottolini, L., Della Ventura, G., Bellatreccia, F. (2008): The crystal chemistry of Li in gadolinite. *Am. Mineral.*, **93**, 996–1004.
- Černá, I., Černý, P., Selway, J.B., Chapman, R. (2002): Paragenesis and origin of secondary berylllophosphates: beryllonite and hydroxylherderite from the Bep granitic pegmatite, southeastern Manitoba, Canada. *Can. Mineral.*, **40**, 1339–1345.

- Černý, P. (1991): Rare-element granitic pegmatites. Part I: anatomy and internal evolution of pegmatite deposits. *Geosci. Canada*, **18**, 49–67.
- Černý, P. (2002): Mineralogy of beryllium in granitic pegmatites. in “Beryllium: mineralogy, petrology, and geochemistry”, Reviews in Mineralogy and Geochemistry, Vol. **50**, E.S. Grew, ed. Mineralogical Society of America, Chantilly, Virginia, 405–444.
- Černý, P. & Ercit, T.S. (2005): The classification of granitic pegmatites revisited. *Can. Mineral.*, **43**, 2005–2026.
- Demartin, F., Pilati, T., Diella, V., Gentile, P., Gramaccioli, C.M. (1993): A crystal-chemical investigation of alpine gadolinite. *Can. Mineral.*, **30**, 127–136.
- Demartin, F., Minaglia, A., Gramaccioli, C.M. (2001): Characterization of gadolinite-group minerals using crystallographic data only: the case of hingganite-(Y) from Cuasso al Monte, Italy. *Can. Mineral.*, **39**, 1105–1114.
- Ding, X., Bai, G., Yuan, Z., Sun, L. (1981): Yttroceberysite, a new Ce–Be-rich silicate. *Geol. Rev.*, **27**, 459–466.
- Ding, X., Bai, G., Yuan, Z., Liu, L. (1984): Hingganite [(Y,Ce)BeSiO₄(OH)]: new data. *Yanshi Kuangwu Ji Ceshi*, **3**, 46–48 (in Chinese).
- Embrey, P.H. & Hey, M.H. (1970): “Type” specimens in mineralogy. *Mineral. Rec.*, **1**, 102–104.
- Foit, F.F. & Gibbs, G.V. (1975): Refinement of $\text{NiYb}_2\text{Be}_2\text{Si}_2\text{O}_{10}$, a gadolinite-type structure. *Z. Kristallogr.*, **141**, 375–386.
- Foit, F.F., Phillips, M.W., Gibbs, G.V. (1973): A refinement of the crystal structure of datolite, $\text{CaBSiO}_4(\text{OH})$. *Am. Mineral.*, **58**, 909–914.
- Foord, E.E., Gaines, R.V., Crock, J.G., Simmons, W.B., Barbosa, C. P. (1986): Minasgeraisite, a new member of the gadolinite group from Minas Gerais, Brazil. *Am. Mineral.*, **71**, 603–607.
- Gibson, S.J. & Ehlmann, A.J. (1970): Annealing characteristics of metamict gadolinite from Rode Ranch, Texas. *Am. Mineral.*, **55**, 288–291.
- Giles, W.B. (1903): Bakerite (a new borosilicate of calcium) and howlite from California. *Mineral. Mag.*, **13**, 353–355.
- Grew, E.S. (1996): Borosilicates (exclusive of tourmaline) and boron in rock-forming minerals in metamorphic environments. in “Boron: mineralogy, petrology and geochemistry”, Reviews in Mineralogy and Geochemistry, Vol. **33**, E.S. Grew & L.M. Anowitz, eds. Mineralogical Society of America, Chantilly, Virginia, 387–502.
- Grew, E.S. (2002): Mineralogy, petrology and geochemistry of beryllium: an introduction and list of beryllium minerals. in “Beryllium: mineralogy, petrology, and geochemistry”, Reviews in Mineralogy and Geochemistry, Vol. **50**, E.S. Grew, ed. Mineralogical Society of America, Chantilly, Virginia, 1–76.
- Grew, E.S., Locock, A.J., Mills, S.J., Galuskina, I.O., Galuskin, E. V., Hålenius, U. (2013): Nomenclature of the garnet supergroup. *Am. Mineral.*, **98**, 785–811.
- Haidinger, W. (1828): On herderite, a new mineral species. *Philos. Mag. Ann. Philos. New Ser.*, **4**, 1–3.
- Hansen, S., Fälth, L., Johnsen, O. (1984a): Bergslagite, a mineral with tetrahedral beryllloarsenate sheet anions. *Z. Kristallogr.*, **160**, 73–80.
- Hansen, S., Fälth, L., Petersen, O.V., Johnsen, O. (1984b): Bergslagite, a new mineral species from Långban, Sweden. *Neues Jb. Miner. Mh.*, **1984**, 257–262.
- Harlow, G.G. & Hawthorne, F.C. (2008): Herderite from Mogok, Myanmar, and comparison with hydroxyl-herderite from Ehrenfriedersdorf, Germany. *Am. Mineral.*, **93**, 1545–1549.

985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045

- 1046 Hatert, F. & Burke, E.A.J. (2008): The IMA-CNMNC dominant-
1047 constituent rule revisited and extended. *Can. Mineral.*, **46**,
1048 717–728.
- 1049 Q3 Hirtopanu, P., Uduba, A.G., Scott, P. (2003): Minerals of the
1050 metamorphosed Mn–Fe deposits in Romania: old deposits, new
1051 species. *Acta Mineral. Petrogr. Abstr. Ser.*, 44.
- 1052 Holtstam, D. & Andersson, U.B. (2007): The REE minerals of the
1053 Bastnäs-type deposits, south-central Sweden. *Can. Mineral.*, **45**,
1054 1073–1114.
- 1055 Ito, J. (1967): Synthesis of calciogadolinite. *Am. Mineral.*, **52**,
1056 1523–1527.
- 1057 Ito, J. & Hafner, S.S. (1974): Synthesis and study of gadolinites. *Am.*
1058 *Mineral.*, **59**, 700–708.
- 1059 Ito, T. & Mori, H. (1953): The crystal structure of datolite. *Acta*
1060 *Crystallogr.*, **6**, 24–32.
- 1061 Jambor, J.L., Roberts, A.C., Grice, J.D., Birkett, T.C., Groat, L.A.,
1062 Zajac, S. (1998): Gerenite-(Y), $(\text{Ca,Na})_2(\text{Y,REE})_3\text{Si}_6\text{O}_{18}\cdot 2\text{H}_2\text{O}$,
1063 a new mineral species, and an associated Y-bearing gadolinite-
1064 group mineral, from the Strange Lake peralkaline complex,
1065 Québec-Labrador. *Can. Mineral.*, **36**, 793–800.
- 1066 King, G.S.D. & Sengier-Roberts, L. (1988): Drugmanite,
1067 $\text{Pb}_2(\text{Fe}_{0.78}\text{Al}_{0.22})\text{H}(\text{PO}_4)_2(\text{OH})_2$: its crystal structure and place
1068 in the datolite group. *Bull. Minéral.*, **111**, 431–437.
- 1069 Klaproth, M.H. (1802): Beiträge zur chemischen Kenntniss der
1070 Mineralkörper 3. Rottmann, Berlin, 52.
- 1071 — (1806): Chemische Untersuchung des Datoliths. *Neues Allg. J.*
1072 *Chem.*, **6**, 107–110.
- 1073 Kozłowski, A. & Dzierżanowski, P. (2007): Gadolinite from the
1074 Michalowice quarry, Karkonosze Massif, SW Poland. *Mineral.*
1075 *Pol.*, **31**, 185–188.
- 1076 Kusachi, I., Henmi, Ch., Kobayashi, Sh. (1994): Chemical
1077 composition of bakerite from Fuka, Okayama Prefecture,
1078 Japan. *Mineral. J.*, **17**, 111–117.
- 1079 Kutzschbach, M., Wunder, B., Rhede, D., Koch-Müller, M., Ertl, A.,
1080 Giester, G., Heinrich, W., Franz, G. (2016): Tetrahedral boron in
1081 natural and synthetic HP/UHP tourmaline: evidence from
1082 Raman spectroscopy, EMPA, and single crystal XRD. *Am.*
1083 *Mineral.*, **101**, 93–104.
- 1084 Leavens, P.B., Dunn, P.J., Gaines, R.V. (1978): Compositional and
1085 refractive index variations of the herderite-hydroxyl-herderite
1086 series. *Am. Mineral.*, **63**, 913–917.
- 1087 Levinson, A.A. (1966): A system of nomenclature for rare-earth
1088 minerals. *Am. Mineral.*, **51**, 152–158.
- 1089 Lulu, X.M. & Zhi-Zhong, P. (1986): Crystal structure of xinganite.
1090 *Geochemistry*, **5**, 280–285.
- 1091 Majka, J., Pršek, J., Budzyn, B., Bačík, P., Barker, A.K., Łodzinski,
1092 M. (2011): Fluorapatite-hingganite-(Y) coronas as products of
1093 fluid induced xenotime-(Y) breakdown in the Skoddefjellet
1094 pegmatite, Svalbard. *Mineral. Mag.*, **75**, 159–167.
- 1095 Massaneck, A., Rank, K., Weber, W. (1999): Die mineralogischen
1096 Sammlungen des Abraham Gottlob Werner. *Lapis*, **24**, 21–31.
- 1097 McGee, J.J. & Anovitz, L.M. (1996): Electron probe microanalysis
1098 of geologic materials for boron. *Rev. Mineral.*, **33**, 771–788.
- 1099 McGee, J.J., Slack, J.F., Herrington, C.R. (1991): Boron analysis by
1100 electron microprobe using MoB_4C layered synthetic crystals.
1101 *Am. Mineral.*, **76**, 681–684.
- 1102 Miyawaki, R. & Nakai, I. (1996): Crystal chemical aspects of rare
1103 earth minerals. in “Rare earth minerals. Chemistry, origin and
1104 ore deposits”, The Mineralogical Society Series, Vol. 7, A.P.
1105 Jones, F. Wall, C.T. Williams, eds. Chapman & Hall, London,
1106 21–40.
- Miyawaki, R., Nakai, I., Nagashima, K. (1984): A refinement of the
crystal structure of gadolinite. *Am. Mineral.*, **69**, 948–953.
- , —, — (1985): Structure of homilite, $\text{Ca}_{2.00}(\text{Fe}_{0.90}\text{Mn}_{0.03})$
 $\text{B}_{2.00}\text{Si}_{2.00}\text{O}_{9.86}(\text{OH})_{0.14}$. *Acta Crystallogr.*, **C41**, 13–15.
- Miyawaki, R., Nakai, I., Nagashima, K., Okamoto, A., Isobe, T.
(1987): The first occurrence of hingganite, hellandite and
wodginite in Japan. *J. Mineral. Soc. Jpn. (Kobutsugaku Zasshi)*,
18, 17–30 (in Japanese with English abstract).
- Miyawaki, R., Matsubara, S., Yokoyama, K., Okamoto, A. (2007):
Hingganite-(Ce) and hingganite-(Y) from Tahara, Hirukawa-
mura, Gifu Prefecture, Japan: the description on a new mineral
species of the Ce-analogue of hingganite-(Y) with a refinement
of the crystal structure of hingganite-(Y). *J. Mineral. Petrol.*
Sci., **102**, 1–7.
- Momma, K. & Izumi, F. (2011): VESTA 3 for three-dimensional
visualization of crystal, volumetric and morphology data. *J.*
Appl. Crystallogr., **44**, 1272–1276.
- Mrose, M.E. & von Knorring, O. (1959): The mineralogy of
väyrynenite, $(\text{Mn,Fe})\text{Be}(\text{PO}_4)(\text{OH})^*$. *Z. Kristallogr.*, **112**, 275–
288.
- Nakai, T. (1938): On calciogadolinite, a new variety of gadolinite
found in Tadachi, Nagano. *Bull. Chem. Soc. Jpn.*, **13**, 591–594.
- Ottolini, L., Cámara, F., Hawthorne, F.C., Stirling, J. (2002): SIMS
matrix effects in the analysis of light elements in silicate
minerals: comparison with SREF and EMPA data. *Am.*
Mineral., **87**, 1477–1485.
- Paijkull, S.R. (1876): Homilit, ett mineral från Brevig i Norge. *Geol.*
Fören. Stock. För., **3**, 229–232.
- Palache, C., Berman, H., Frondel, C. (1951): Dana’s system of Q4
mineralogy, 7th edn., Vol. II, 363.
- Pasero, M., Kampf, A.R., Ferraris, C., Pekov, I.V., Rakovan, J.,
White, T.J. (2010): Nomenclature of the apatite supergroup
minerals. *Eur. J. Mineral.*, **22**, 163–179.
- Pekov, I.V., Voloshin, A.V., Pushcharovskii, D.Yu., Rastsvetaeva, R.
K., Chukanov, N.V., Belakovskii, D.I. (2000): New data on
calcybeborosilite-(Y) $(\text{REE,Ca})_2(\text{B,Be})_2(\text{SiO}_4)_2(\text{OH,O})_2$. *Vestn.*
Mosk. Univ. Geol., **55**, 62–70.
- Pemberton, H.E. (1971): Type locality for bakerite. *Am. Mineral.*,
56, 1109–1110.
- Penfield, S.L. (1894): On the crystallization of herderite. *Am. J. Sci.*,
147, 329–339.
- Perchiazzi, N., Gualtieri, A.F., Merlino, S., Kampf, A.R. (2004): The
atomic structure of bakerite and its relationship to datolite. *Am.*
Mineral., **89**, 767–776.
- Pezzotta, F., Diella, V., Guastoni, A. (1999): Chemical and
paragenetic data on gadolinite-group minerals from Baveno
and Cuasso al Monte, southern Alps, Italy. *Am. Mineral.*, **84**,
782–789.
- Pierini, G. (2004): Homilite in un proietto vulcanico a Tre Croci di
Vetralla (VT). *Micro (notizie mineralogiche)*, **2004**, 5–12.
- Povarennykh, A.S. & Dusmatov, V.D. (1970): Infrared absorption
spectra of new minerals from alkaline pegmatites of central
Asia. *Konst. Svoistva Mineralov, Akad. Nauk Ukr. S.S.R.,*
Respub. Mezhvedom. Sb., **4**, 3–9.
- Pršek, J., Ondrejka, M., Bačík, P., Budzyn, B., Uher, P. (2010):
Metamorphic-hydrothermal REE minerals in the Bacúch
magnetite deposit, Western Carpathians, Slovakia: (Sr,S)-rich
monazite-(Ce) and Nd-dominant hingganite. *Can. Mineral.*, **48**,
81–94.
- Raade, G., Kolitsch, U., Husdal, T.A. (2006): Si-rich bergslagite from
a granitic pegmatite at Tennvatn, north Norway. *GFF*, **128**, 65–68.

- 1168 Rinaldi, R., Gatta, G.D., Angel, R.J. (2010): Crystal chemistry and
1169 low-temperature behavior of datolite: a single-crystal X-ray
1170 diffraction study. *Am. Mineral.*, **95**, 1413–1421. 1197
- 1171 Sabina, A.P. (1978): Some new mineral occurrences in Canada. 1198
1172 Geological Survey of Canada, Paper 78-1A, 253–258. 1199
- 1173 Schnorrer, G. & Schäfer, H. (1999): Drugmanit, $\text{Pb}_2(\text{Fe}^{3+}, \text{Al})[\text{OH}/$
1174 $\text{PO}_4]_2 \cdot \text{H}_2\text{O}$, ein drittes Vorkommen aus der Eifel (Grube Neue
1175 Hoffnung). *Der Aufschluss*, **50**, 225–228. 1200
- 1176 Segalstad, T.V. & Larsen, A.O. (1978): Gadolinite-(Ce) from Skien,
1177 southern Oslo region, Norway. *Am. Mineral.*, **63**, 188–195. 1201
- 1178 Semenov, E.I., Dusmatov, V.D., Samsonova, N.S. (1963): Yttrium-
1179 beryllium minerals of the datolite group. *Kristallografiya*, **8**,
1180 667–679. 1202
- 1181 Shannon, R.D. (1976): Revised effective ionic radii and systematic
1182 studies of interatomic distances in halides and chalcogenides. 1203
1183 *Acta Crystallogr.*, **A32**, 751–767. 1204
- 1184 Škoda, R., Plášil, J., Čopjaková, R., Novák, M., Jonsson, E.,
1185 Vašinová Galiová, M., Holtstam, D. (2016): Gadolinite-(Nd),
1186 IMA 2016-013. CNMNC Newsletter No. 32, August 2016, page
1187 916. *Mineral. Mag.*, **80**, 915–922. 1205
- 1188 Uher, P., Ondrejka, M., Konečný, P. (2009): Magmatic and post-
1189 magmatic Y-REE-Th phosphate, silicate and Nb-Ta-Y-REE
1190 oxide minerals in A-type metagranite: an example from the
1191 Turčok massif, the Western Carpathians, Slovakia. *Mineral.*
1192 *Mag.*, **73**, 1009–1025. 1206
- 1193 Uher, P., Bačík, P., Ozdín, D., Hornáček, M. (2010): Datolite from
1194 Permian metabasalts in Lošonec (Malé Karpaty Mts., SW
1195 Slovakia). *Bull. Mineral. Petrol. Odd. Nár. Muz. (Praha)*, **18**,
1196 91–95 (in Slovak). 1207
- 1208 Van Tassel, R., Fransolet, A.-M., Abraham, K. (1979): Drugmanite,
1209 $\text{Pb}_2(\text{Fe}^{3+}, \text{Al})(\text{PO}_4)_2(\text{OH}) \cdot \text{H}_2\text{O}$, a new mineral from Richelle,
1210 Belgium. *Mineral. Mag.*, **43**, 463–467. 1211
- 1212 Volborth, J. (1954): Väyryneniiti, $\text{BeMnPO}_4(\text{OH}, \text{F})$. *Geologi*, **6**, 7. 1212
- 1213 Voloshin, A.V., Pakhomovskii, J.A., Menshikov, I.P., Povarennykh,
1214 A.S., Matvienko, E.N., Iakubovich, O.V. (1983): Hingganite
1215 (Yb), a new mineral from the amazonite pegmatites of the
1216 Kola-peninsula. *Dokl. Akad. Nauk SSSR*, **270**, 1188–1192. 1217
- 1218 Voloshin, A.V., Pakhomovskii, J.A., Sorokhtina, N.V. (2002):
1219 Chemical composition of the gadolinite-group minerals from
1220 the amazonite pegmatites in the Kola Peninsula. *Vestnik*
1221 *Murmanskiy Gosudarstvennyy Tekhnicheskii Universitet*, **5**,
1222 61–70 (in Russian). 1223
- 1224 Yakubovich, O.V., Matvinenko, E.N., Voloshin, A.V., Simonov, M.
1225 A. (1983): The crystal structure of hingganite-(Yb),
1226 $(\text{Y}_{0.51}\text{TR}_{0.36}\text{Ca}_{0.13})\text{Fe}_{0.065}\text{Be}[\text{SiO}_4](\text{OH})$. *Kristallografiya*, **28**,
1227 457–460. 1228
- 1229 Zaccarini, F., Morales-Ruano, S., Scacchetti, M., Garuti, G., Heide,
1230 K. (2008): Investigation of datolite $\text{CaBSiO}_4(\text{OH})$ from basalts
1231 in the Northern Apennines ophiolites (Italy): genetic implica-
1232 tions. *Chem. Erde*, **68**, 265–277. 1233
- 1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250

Received 11 October 2016

Modified version received 23 January 2017

Accepted 23 January 2017

uncorrected proof